



Q -polynomial distance-regular graphs with $a_1 = 0$ and $a_2 \neq 0$

Štefko Miklavič

Primorska Institute for Natural Science and Technology, University of Primorska, 6000 Koper, Slovenia

Received 7 August 2007; accepted 1 February 2008

Available online 14 March 2008

Abstract

Let Γ denote a Q -polynomial distance-regular graph with diameter $D \geq 3$ and intersection numbers $a_1 = 0$, $a_2 \neq 0$. Let X denote the vertex set of Γ and let $A \in \text{Mat}_X(\mathbb{C})$ denote the adjacency matrix of Γ . Fix $x \in X$ and let $A^* \in \text{Mat}_X(\mathbb{C})$ denote the corresponding dual adjacency matrix. Let T denote the subalgebra of $\text{Mat}_X(\mathbb{C})$ generated by A , A^* . We call T the *Terwilliger algebra* of Γ with respect to x . We show that up to isomorphism there exists a unique irreducible T -module W with endpoint 1. We show that W has dimension $2D - 2$. We display a basis for W which consists of eigenvectors for A^* . We display the action of A on this basis. We show that W appears in the standard module of Γ with multiplicity $k - 1$, where k is the valency of Γ .

© 2008 Published by Elsevier Ltd

1. Introduction

Let Γ denote a Q -polynomial distance-regular graph with diameter $D \geq 3$ and intersection numbers a_i , b_i , c_i (see Section 2 for formal definitions). We recall the Terwilliger algebra of Γ . Let X denote the vertex set of Γ and let $A \in \text{Mat}_X(\mathbb{C})$ denote the adjacency matrix of Γ . Fix a “base vertex” $x \in X$ and let $A^* \in \text{Mat}_X(\mathbb{C})$ denote the corresponding dual adjacency matrix. Let $T = T(x)$ denote the subalgebra of $\text{Mat}_X(\mathbb{C})$ generated by A , A^* . The algebra T is called the *Terwilliger algebra* of Γ with respect to x [26]. T is closed under the conjugate-transpose map so T is semi-simple [26, Lemma 3.4(i)]. Therefore each T -module is a direct sum of irreducible T -modules. Describing the irreducible T -modules is an active area of research [3–17, 25–27].

In this description there is an important parameter called the *endpoint* which we now recall. Let W denote an irreducible T -module. By the *endpoint* of W we mean $\min\{i \mid 0 \leq i \leq$

E-mail address: stefko.miklavic@upr.si.

D , $E_i^*W \neq 0$), where $E_i^* \in \text{Mat}_X(\mathbb{C})$ is the projection onto the i th subconstituent of Γ with respect to x [26, p. 378]. There exists a unique irreducible T -module with endpoint 0 [12, Proposition 8.4]; for a detailed description see [7,12]. In this paper we consider irreducible T -modules with endpoint 1.

We are going to make an assumption about the intersection numbers of Γ . In order to motivate things let us review the situation for $a_1 = 0$. In this case either (i) $a_i = 0$ for $0 \leq i \leq D$ (the bipartite case); or (ii) $a_i = 0$ for $0 \leq i \leq D - 1$ and $a_D \neq 0$ (the almost bipartite case); or (iii) $a_i \neq 0$ for $2 \leq i \leq D$ [21, Theorem 6.3]. In case (i) (resp. case (ii)) the irreducible T -modules were determined in [7–9] (resp. [4,6]). In this paper we assume case (iii) and describe the irreducible T -modules with endpoint 1. Our results are summarized as follows. We show that up to isomorphism there exists a unique irreducible T -module W with endpoint 1. We show that W has dimension $2D - 2$. We display a basis for W which consists of eigenvectors of A^* . We display the action of A on this basis. We show that W appears in the standard module of Γ with multiplicity $k - 1$, where k is the valency of Γ . We remark that the following Q -polynomial distance-regular graphs satisfy $a_1 = 0$ and $a_i \neq 0$ for $2 \leq i \leq D$: the Hermitian forms graphs with $r = 2$ [2, Section 9.5.C] and the Witt graph M_{23} [2, Section 11.4.B]. We discuss these examples later in the paper.

2. Preliminaries

In this section we review some definitions and basic results concerning distance-regular graphs. See the book of Brouwer, Cohen and Neumaier [2] for more background information.

Let \mathbb{C} denote the complex number field and let X denote a nonempty finite set. Let $\text{Mat}_X(\mathbb{C})$ denote the \mathbb{C} -algebra consisting of all matrices whose rows and columns are indexed by X and whose entries are in \mathbb{C} . Let $V = \mathbb{C}^X$ denote the vector space over \mathbb{C} consisting of column vectors whose coordinates are indexed by X and whose entries are in \mathbb{C} . We observe that $\text{Mat}_X(\mathbb{C})$ acts on V by left multiplication. We call V the *standard module*. We endow V with the Hermitian inner product $\langle \cdot, \cdot \rangle$ that satisfies $\langle u, v \rangle = u^t \bar{v}$ for $u, v \in V$, where t denotes transpose and $\bar{\cdot}$ denotes complex conjugation. For $y \in X$ let \hat{y} denote the element of V with a 1 in the y coordinate and 0 in all other coordinates. We observe that $\{\hat{y} \mid y \in X\}$ is an orthonormal basis for V . The following will be useful: for each $B \in \text{Mat}_X(\mathbb{C})$ we have

$$\langle u, Bv \rangle = \langle \bar{B}^t u, v \rangle \quad (u, v \in V). \quad (1)$$

Let $\Gamma = (X, R)$ denote a finite, undirected, connected graph, without loops or multiple edges, with vertex set X and edge set R . Let ∂ denote the path-length distance function for Γ , and set $D := \max\{\partial(x, y) \mid x, y \in X\}$. We call D the *diameter* of Γ . For a vertex $x \in X$ and an integer $i \geq 0$ let $\Gamma_i(x)$ denote the set of vertices at distance i from x . We use the abbreviation $\Gamma(x) = \Gamma_1(x)$. For an integer $k \geq 0$ we say that Γ is *regular with valency k* whenever $|\Gamma(x)| = k$ for all $x \in X$. We say that Γ is *distance-regular* whenever for all integers h, i, j ($0 \leq h, i, j \leq D$) and for all vertices $x, y \in X$ with $\partial(x, y) = h$, the number

$$p_{ij}^h = |\Gamma_i(x) \cap \Gamma_j(y)|$$

is independent of x and y . The p_{ij}^h are called the *intersection numbers* of Γ .

For the rest of this paper we assume that Γ is distance-regular with diameter $D \geq 3$. Note that $p_{ij}^h = p_{ji}^h$ for $0 \leq h, i, j \leq D$. For convenience set $c_i := p_{1,i-1}^1$ ($1 \leq i \leq D$), $a_i := p_{1i}^1$ ($0 \leq i \leq D$), $b_i := p_{1,i+1}^1$ ($0 \leq i \leq D - 1$), $k_i := p_{ii}^0$ ($0 \leq i \leq D$), and $c_0 = b_D = 0$.

By the triangle inequality the following hold for $0 \leq h, i, j \leq D$: (i) $p_{ij}^h = 0$ if one of h, i, j is greater than the sum of the other two; (ii) $p_{ij}^h \neq 0$ if one of h, i, j equals the sum of the other two. In particular $c_i \neq 0$ for $1 \leq i \leq D$ and $b_i \neq 0$ for $0 \leq i \leq D-1$. We observe that Γ is regular with valency $k = k_1 = b_0$ and that $c_i + a_i + b_i = k$ for $0 \leq i \leq D$. Note that $k_i = |\Gamma_i(x)|$ for $x \in X$ and $0 \leq i \leq D$. By [2, p. 127],

$$k_i = \frac{b_0 b_1 \cdots b_{i-1}}{c_1 c_2 \cdots c_i} \quad (0 \leq i \leq D). \quad (2)$$

We recall the Bose–Mesner algebra of Γ . For $0 \leq i \leq D$ let A_i denote the matrix in $\text{Mat}_X(\mathbb{C})$ with (x, y) -entry

$$(A_i)_{xy} = \begin{cases} 1 & \text{if } \partial(x, y) = i, \\ 0 & \text{if } \partial(x, y) \neq i \end{cases} \quad (x, y \in X). \quad (3)$$

We call A_i the i th *distance matrix* of Γ . We use the abbreviation $A := A_1$ and call this the *adjacency matrix* of Γ . We observe that (ai) $A_0 = I$; (aii) $\sum_{i=0}^D A_i = J$; (aiii) $\overline{A_i} = A_i$ ($0 \leq i \leq D$); (aiv) $A_i^t = A_i$ ($0 \leq i \leq D$); (av) $A_i A_j = \sum_{h=0}^D p_{ij}^h A_h$ ($0 \leq i, j \leq D$), where I (resp. J) denotes the identity matrix (resp. all 1's matrix) in $\text{Mat}_X(\mathbb{C})$. Using these facts we find that A_0, A_1, \dots, A_D is a basis for a commutative subalgebra M of $\text{Mat}_X(\mathbb{C})$. We call M the *Bose–Mesner algebra* of Γ . It turns out that A generates M [1, p. 190]. By [2, p. 45], M has a second basis E_0, E_1, \dots, E_D such that (ei) $E_0 = |X|^{-1} J$; (eii) $\sum_{i=0}^D E_i = I$; (eiii) $\overline{E_i} = E_i$ ($0 \leq i \leq D$); (eiv) $E_i^t = E_i$ ($0 \leq i \leq D$); (ev) $E_i E_j = \delta_{ij} E_i$ ($0 \leq i, j \leq D$). We call E_0, E_1, \dots, E_D the *primitive idempotents* of Γ .

We now recall the Krein parameters. Let \circ denote the entrywise product in $\text{Mat}_X(\mathbb{C})$. Observe that $A_i \circ A_j = \delta_{ij} A_i$ for $0 \leq i, j \leq D$, so M is closed under \circ . Thus there exist complex scalars q_{ij}^h ($0 \leq h, i, j \leq D$) such that

$$E_i \circ E_j = |X|^{-1} \sum_{h=0}^D q_{ij}^h E_h \quad (0 \leq i, j \leq D).$$

By [2, Proposition 4.1.5], q_{ij}^h is real and nonnegative for $0 \leq h, i, j \leq D$. The q_{ij}^h are called the *Krein parameters* of Γ . The graph Γ is said to be *Q-polynomial* (with respect to the given ordering E_0, E_1, \dots, E_D of the primitive idempotents) whenever for $0 \leq h, i, j \leq D$, $q_{ij}^h = 0$ (resp. $q_{ij}^h \neq 0$) whenever one of h, i, j is greater than (resp. equal to) the sum of the other two. From now on assume that Γ is *Q-polynomial* with respect to E_0, E_1, \dots, E_D .

We recall the dual eigenvalues of Γ . Define complex scalars $\theta_0^*, \theta_1^*, \dots, \theta_D^*$ by

$$E_1 = |X|^{-1} \sum_{i=0}^D \theta_i^* A_i.$$

We call the sequence $\theta_0^*, \theta_1^*, \dots, \theta_D^*$ the *dual eigenvalue sequence* of Γ for the given *Q-polynomial* structure. The scalars $\theta_0^*, \theta_1^*, \dots, \theta_D^*$ are real and mutually distinct [26, Lemma 3.11(iii)].

3. The Terwilliger algebra

In this section we recall the dual Bose–Mesner algebra and the Terwilliger algebra of Γ . Fix a vertex $x \in X$. We view x as a “base vertex”. For $0 \leq i \leq D$ let $E_i^* = E_i^*(x)$ denote the diagonal

matrix in $\text{Mat}_X(\mathbb{C})$ with (y, y) -entry

$$(E_i^*)_{yy} = \begin{cases} 1 & \text{if } \partial(x, y) = i, \\ 0 & \text{if } \partial(x, y) \neq i \end{cases} \quad (y \in X). \quad (4)$$

We call E_i^* the i th *dual idempotent* of Γ with respect to x [26, p. 378]. We observe that (i) $\sum_{i=0}^D E_i^* = I$; (ii) $\overline{E_i^*} = E_i^*(0 \leq i \leq D)$; (iii) $E_i^{*t} = E_i^*(0 \leq i \leq D)$; (iv) $E_i^* E_j^* = \delta_{ij} E_i^*(0 \leq i, j \leq D)$. By these facts $E_0^*, E_1^*, \dots, E_D^*$ form a basis for a commutative subalgebra $M^* = M^*(x)$ of $\text{Mat}_X(\mathbb{C})$. We call M^* the *dual Bose–Mesner algebra* of Γ with respect to x [26, p. 378]. For $0 \leq i \leq D$ we have

$$E_i^* V = \text{Span}\{\hat{y} \mid y \in X, \partial(x, y) = i\}$$

so $\dim E_i^* V = k_i$. We call $E_i^* V$ the i th *subconstituent* of Γ with respect to x . Note that

$$V = E_0^* V + E_1^* V + \dots + E_D^* V \quad (\text{orthogonal direct sum}). \quad (5)$$

Moreover E_i^* is the projection from V onto $E_i^* V$ for $0 \leq i \leq D$.

Let $A^* = A^*(x)$ denote the diagonal matrix in $\text{Mat}_X(\mathbb{C})$ with (y, y) -entry

$$A_{yy}^* = |X| E_{xy} \quad (y \in X),$$

where $E = E_1$. By the construction, $A^* = \sum_{i=0}^D \theta_i^* E_i^*$. We call A^* the *dual adjacency matrix* of Γ with respect to x . By [26, Lemma 3.11(ii)], A^* generates M^* .

We recall the Terwilliger algebra of Γ . Let $T = T(x)$ denote the subalgebra of $\text{Mat}_X(\mathbb{C})$ generated by M, M^* . We call T the *Terwilliger algebra* of Γ with respect to x [26, Definition 3.3]. Recall that M (resp. M^*) is generated by A (resp. A^*) so T is generated by A, A^* . We observe that T has finite dimension. By construction T is closed under the conjugate-transpose map so T is semi-simple [26, Lemma 3.4(i)].

By a T -module we mean a subspace W of V such that $BW \subseteq W$ for all $B \in T$. Let W denote a T -module. Then W is said to be *irreducible* whenever W is nonzero and W contains no T -modules other than 0 and W . Assume that W is irreducible. Then A and A^* act on W as a tridiagonal pair [17, Example 1.4]. We refer the reader to [17–20, 22, 23] and the references therein for background on tridiagonal pairs.

By [14, Corollary 6.2] any T -module is an orthogonal direct sum of irreducible T -modules. In particular the standard module V is an orthogonal direct sum of irreducible T -modules. Let W, W' denote T -modules. By an *isomorphism of T -modules* from W to W' we mean an isomorphism of vector spaces $\sigma : W \rightarrow W'$ such that $(\sigma B - B\sigma)W = 0$ for all $B \in T$. The T -modules W, W' are said to be *isomorphic* whenever there exists an isomorphism of T -modules from W to W' . By [7, Lemma 3.3] any two nonisomorphic irreducible T -modules are orthogonal. Let W denote an irreducible T -module. By [26, Lemma 3.4(iii)] W is an orthogonal direct sum of the nonvanishing spaces among $E_0^* W, E_1^* W, \dots, E_D^* W$. By the *endpoint* of W we mean $\min\{i \mid 0 \leq i \leq D, E_i^* W \neq 0\}$. By the *diameter* of W we mean $|\{i \mid 0 \leq i \leq D, E_i^* W \neq 0\}| - 1$.

By [12, Proposition 8.3, Proposition 8.4] $M\hat{x}$ is the unique irreducible T -module with endpoint 0 and the unique irreducible T -module with diameter D . Moreover $M\hat{x}$ is the unique irreducible T -module on which E_0 does not vanish. We call $M\hat{x}$ the *primary module*. From now until the end of Section 10 we adopt the following notational convention.

Notation 3.1. Let $\Gamma = (X, R)$ denote a Q -polynomial distance-regular graph with diameter $D \geq 3$ and valency k . We assume that $a_1 = 0$, $a_2 \neq 0$ and note that $a_i \neq 0$ for $2 \leq i \leq D$ [21, Theorem 6.3]. Let A_0, A_1, \dots, A_D denote the distance matrices of Γ , and let V denote the standard module of Γ . We fix $x \in X$ and let $E_i^* = E_i^*(x)$ ($0 \leq i \leq D$), and $T = T(x)$ denote the corresponding dual idempotents and Terwilliger algebra, respectively. Let $\theta_0^*, \theta_1^*, \dots, \theta_D^*$ denote the dual eigenvalue sequence of Γ for the given Q -polynomial structure.

4. The sets D_j^i

With reference to Notation 3.1, in this section we define certain subsets D_j^i of X and explore their properties.

Definition 4.1. With reference to Notation 3.1 fix $y \in \Gamma(x)$. For $0 \leq i, j \leq D$ we define $D_j^i = D_j^i(x, y)$ by

$$D_j^i = \Gamma_i(x) \cap \Gamma_j(y).$$

For notational convenience we set $D_j^i = \emptyset$ if i or j is contained in $\{-1, D+1\}$.

Lemma 4.2. With reference to Notation 3.1 the following (i), (ii) hold for $0 \leq i, j \leq D$.

- (i) $|D_j^i| = p_{ij}^1$.
- (ii) $D_j^i = \emptyset$ if and only if $p_{ij}^1 = 0$.

Proof. (i) Immediate from the definition of p_{ij}^1 and D_j^i .

(ii) Immediate from (i) above. ■

Lemma 4.3 ([2, p. 134]). With reference to Notation 3.1 the following (i), (ii) hold.

- (i) $p_{i-1,i}^1 = p_{i,i-1}^1 = c_i k_i k^{-1}$ ($1 \leq i \leq D$).
- (ii) $p_{ii}^1 = a_i k_i k^{-1}$ ($0 \leq i \leq D$).

Lemma 4.4. With reference to Notation 3.1 the following (i)–(iii) hold.

- (i) $p_{i-1,i}^1 \neq 0$, $p_{i,i-1}^1 \neq 0$ ($1 \leq i \leq D$).
- (ii) $p_{00}^1 = 0$, $p_{11}^1 = 0$, $p_{ii}^1 \neq 0$ ($2 \leq i \leq D$).
- (iii) $p_{ij}^1 = 0$ if $|i - j| \notin \{0, 1\}$.

Proof. (i), (ii) Immediate from Lemma 4.3.

(iii) Immediate from the triangle inequality. ■

Theorem 4.5 ([21, Theorem 5.2]). With reference to Notation 3.1 fix $y \in \Gamma(x)$ and use the abbreviation $D_j^i = D_j^i(x, y)$ for $0 \leq i, j \leq D$. Then the following (i), (ii) hold.

- (i) For $1 \leq i \leq D$ each $z \in D_{i-1}^i$ (resp. D_i^{i-1}) is adjacent to

- (a) precisely ρ_i vertices in D_{i-1}^{i-1} ,
- (b) precisely c_{i-1} vertices in D_{i-2}^{i-1} (resp. D_{i-1}^{i-2}),
- (c) precisely $c_i - c_{i-1} - \rho_i$ vertices in D_i^{i-1} (resp. D_{i-1}^i),
- (d) precisely $a_{i-1} - \rho_i$ vertices in D_{i-1}^i (resp. D_i^{i-1}),
- (e) precisely b_i vertices in D_i^{i+1} (resp. D_{i+1}^i),
- (f) precisely $a_i - a_{i-1} + \rho_i$ vertices in D_i^i ,

where

$$\rho_i = a_{i-1} \frac{(\theta_0^* - \theta_{i-1}^*)(\theta_2^* - \theta_i^*) - (\theta_1^* - \theta_{i-1}^*)(\theta_1^* - \theta_i^*)}{(\theta_0^* - \theta_{i-1}^*)(\theta_{i-1}^* - \theta_i^*)} \quad (2 \leq i \leq D), \quad \rho_1 = 0.$$

(ii) For $2 \leq i \leq D$ each $z \in D_i^i$ is adjacent to

- | | | |
|---------------|---------------------------------------|-------------------------------|
| (a) precisely | σ_i | vertices in D_{i-1}^{i-1} , |
| (b) precisely | τ_i | vertices in D_{i+1}^{i+1} , |
| (c) precisely | $c_i - \sigma_i$ | vertices in D_{i-1}^i , |
| (d) precisely | $c_i - \sigma_i$ | vertices in D_i^{i-1} , |
| (e) precisely | $b_i - \tau_i$ | vertices in D_i^{i+1} , |
| (f) precisely | $b_i - \tau_i$ | vertices in D_{i+1}^i , |
| (g) precisely | $a_i - b_i - c_i + \sigma_i + \tau_i$ | vertices in D_i^i , |

where

$$\sigma_i = c_i \frac{(\theta_0^* - \theta_i^*)(\theta_2^* - \theta_i^*) - (\theta_1^* - \theta_{i-1}^*)(\theta_1^* - \theta_i^*)}{(\theta_0^* - \theta_i^*)(\theta_{i-1}^* - \theta_i^*)},$$

$$\tau_i = b_i \frac{(\theta_0^* - \theta_i^*)(\theta_2^* - \theta_i^*) - (\theta_1^* - \theta_i^*)(\theta_1^* - \theta_{i+1}^*)}{(\theta_0^* - \theta_i^*)(\theta_{i+1}^* - \theta_i^*)} \quad (2 \leq i \leq D-1), \quad \tau_D = 0.$$

5. Some products in T

With reference to [Notation 3.1](#), in this section we evaluate several products in T which we shall need later.

Lemma 5.1. With reference to [Notation 3.1](#), for $0 \leq h, i, j \leq D$ and $y, z \in X$ the (y, z) -entry of $E_h^* A_i E_j^*$ is 1 if $\partial(x, y) = h$, $\partial(y, z) = i$, $\partial(x, z) = j$, and 0 otherwise.

Proof. Compute the (y, z) -entry of $E_h^* A_i E_j^*$ by matrix multiplication and simplify the result using (3) and (4). ■

Corollary 5.2 ([26, Lemma 3.2]). With reference to [Notation 3.1](#),

$$E_h^* A_i E_j^* = 0 \quad \text{if and only if } p_{ij}^h = 0 \quad (0 \leq h, i, j \leq D).$$

Proof. Immediate from [Lemma 5.1](#). ■

Corollary 5.3. With reference to [Notation 3.1](#) the following (i)–(iii) hold.

- (i) $E_1^* A E_1^* = 0$.
- (ii) $E_i^* A_{i-1} E_1^* + E_i^* A_i E_1^* + E_i^* A_{i+1} E_1^* = E_i^* J E_1^*$ for $1 \leq i \leq D-1$.
- (iii) $E_D^* A_{D-1} E_1^* + E_D^* A_D E_1^* = E_D^* J E_1^*$.

Proof. (i) By [Corollary 5.2](#) and since $p_{11}^1 = 0$ by [Lemma 4.4\(ii\)](#).

(ii), (iii) For each equation evaluate the right-hand side using assertion (aii) below line (3), and simplify the result using [Corollary 5.2](#) and assertion (i) above line (2). ■

Lemma 5.4. With reference to [Notation 3.1](#), for $0 \leq h, i, j, r, s \leq D$ and $y, z \in X$ the (y, z) -entry of $E_h^* A_r E_i^* A_s E_j^*$ is $|\Gamma_i(x) \cap \Gamma_r(y) \cap \Gamma_s(z)|$ if $\partial(x, y) = h$, $\partial(x, z) = j$, and 0 otherwise.

Proof. Compute the (y, z) -entry of $E_h^* A_r E_i^* A_s E_j^*$ by matrix multiplication and simplify the result using (3) and (4). ■

6. The matrices L, F, R

With reference to [Notation 3.1](#), in this section we recall the matrices L, F, R and use them to interpret [Theorem 4.5](#).

Definition 6.1. With reference to [Notation 3.1](#) we define matrices $L = L(x), F = F(x), R = R(x)$ by

$$L = \sum_{h=1}^D E_{h-1}^* A E_h^*, \quad F = \sum_{h=0}^D E_h^* A E_h^*, \quad R = \sum_{h=0}^{D-1} E_{h+1}^* A E_h^*.$$

Note that $A = L + F + R$ [[7](#), Lemma 4.4]. We call L, F , and R the *lowering matrix*, the *flat matrix*, and the *raising matrix* of Γ with respect to x , respectively.

Lemma 6.2. With reference to [Notation 3.1](#) and [Definition 6.1](#) the following (i)–(v) hold.

- (i) $LE_1^* = E_0^* A E_1^*$.
- (ii) $LE_2^* A E_1^* = b_1 E_1^* + (c_2 - 1) E_1^* A_2 E_1^*$.
- (iii) $LE_i^* A_{i-1} E_1^* = b_{i-1} E_{i-1}^* A_{i-2} E_1^* + (b_{i-1} - \tau_{i-1}) E_{i-1}^* A_{i-1} E_1^* + (c_i - c_{i-1} - \rho_i) E_{i-1}^* A_i E_1^*$ for $3 \leq i \leq D$.
- (iv) $LE_2^* A_2 E_1^* = a_2 E_1^* A_2 E_1^*$.
- (v) $LE_i^* A_i E_1^* = \tau_{i-1} E_{i-1}^* A_{i-1} E_1^* + (a_i - a_{i-1} + \rho_i) E_{i-1}^* A_i E_1^*$ for $3 \leq i \leq D$.

Proof. For each equation and for $z, w \in X$ compute the (z, w) -entry of each side and interpret the results using [Lemmas 5.1](#) and [5.4](#) and [Theorem 4.5](#). ■

Lemma 6.3. With reference to [Notation 3.1](#) and [Definition 6.1](#) the following (i)–(iv) hold.

- (i) $FE_1^* = 0$.
- (ii) $FE_i^* A_{i-1} E_1^* = (a_{i-1} - \rho_i) E_i^* A_{i-1} E_1^* + (c_i - \sigma_i) E_i^* A_i E_1^*$ for $2 \leq i \leq D$.
- (iii) $FE_i^* A_i E_1^* = (a_i - a_{i-1} + \rho_i) E_i^* A_{i-1} E_1^* + (a_i - b_i - c_i + \sigma_i + \tau_i) E_i^* A_i E_1^* + \rho_{i+1} E_i^* A_{i+1} E_1^*$ for $2 \leq i \leq D - 1$.
- (iv) $FE_D^* A_D E_1^* = (a_D - a_{D-1} + \rho_D) E_D^* A_{D-1} E_1^* + (a_D - c_D + \sigma_D) E_D^* A_D E_1^*$.

Proof. For each equation and for $z, w \in X$ compute the (z, w) -entry of each side and interpret the results using [Lemmas 5.1](#) and [5.4](#) and [Theorem 4.5](#). ■

Lemma 6.4. With reference to [Notation 3.1](#) and [Definition 6.1](#) the following (i)–(iv) hold.

- (i) $RE_i^* A_{i-1} E_1^* = c_i E_{i+1}^* A_i E_1^*$ for $1 \leq i \leq D - 1$.
- (ii) $RE_D^* A_{D-1} E_1^* = 0$.
- (iii) $RE_i^* A_i E_1^* = \rho_{i+1} E_{i+1}^* A_i E_1^* + \sigma_{i+1} E_{i+1}^* A_{i+1} E_1^*$ for $2 \leq i \leq D - 1$.
- (iv) $RE_D^* A_D E_1^* = 0$.

Proof. For each equation and for $z, w \in X$ compute the (z, w) -entry of each side and interpret the results using [Lemmas 5.1](#) and [5.4](#) and [Theorem 4.5](#). ■

7. More products in T

With reference to [Notation 3.1](#), in this section we evaluate more products in T which we will need later.

Lemma 7.1. *With reference to [Notation 3.1](#), for $y, z \in \Gamma(x)$ and for $1 \leq i \leq D$ the number $|\Gamma_i(x) \cap \Gamma_{i-1}(y) \cap \Gamma_{i-1}(z)|$ is equal to $c_i k_i k^{-1}$ if $y = z$ and $c_i(c_i - 1)k_i k^{-1} b_1^{-1}$ if $y \neq z$.*

Proof. Assume that $y \neq z$; otherwise the result follows by [Lemma 4.3\(i\)](#). Use the abbreviation $D_j^\ell = D_j^\ell(x, y)$ ($0 \leq \ell, j \leq D$) and note that $z \in D_2^1$. It follows from [Theorem 4.5](#) that the number of paths of length $i - 1$ between z and D_{i-1}^i is independent of x, y, z . Moreover, between any two vertices of Γ which are at distance $i - 1$, there exist exactly $c_1 c_2 \cdots c_{i-1}$ paths of length $i - 1$, implying that $|D_{i-1}^i \cap \Gamma_{i-1}(z)|$ is independent of x, y, z . For $v \in D_{i-1}^i$ we have $|\Gamma_{i-1}(v) \cap \Gamma(x)| = c_i$ so $|\Gamma_{i-1}(v) \cap D_2^1| = c_i - 1$. Counting the number of pairs (w, v) such that $w \in D_2^1, v \in D_{i-1}^i, \partial(w, v) = i - 1$ in two different ways and using the above comments we obtain $|D_{i-1}^i \cap \Gamma_{i-1}(z)| |D_2^1| = |D_{i-1}^i| (c_i - 1)$. The result now follows by [Lemma 4.2\(i\)](#) and [Lemma 4.3\(i\)](#). ■

Corollary 7.2. *With reference to [Notation 3.1](#), for $1 \leq i \leq D$ we have*

$$E_1^* A_{i-1} E_i^* A_{i-1} E_1^* = c_i k_i k^{-1} E_1^* + c_i(c_i - 1) k_i k^{-1} b_1^{-1} E_1^* A_2 E_1^*.$$

Proof. For $z, w \in X$ we show that the (z, w) -entries of both sides are equal. If $z \notin \Gamma(x)$ or $w \notin \Gamma(x)$ then the (z, w) -entry of each side is 0. If $z, w \in \Gamma(x)$ then the (z, w) -entries of both sides are equal by [Lemmas 5.1, 5.4](#) and [7.1](#). The result follows. ■

Lemma 7.3. *With reference to [Notation 3.1](#), for $y, z \in \Gamma(x)$ and for $1 \leq i \leq D$ the number $|\Gamma_i(x) \cap \Gamma_i(y) \cap \Gamma_{i-1}(z)|$ is equal to 0 if $y = z$ and $a_i c_i k_i k^{-1} b_1^{-1}$ if $y \neq z$.*

Proof. Assume that $y \neq z$; otherwise the result is clear. Use the abbreviation $D_j^\ell = D_j^\ell(x, y)$ ($0 \leq \ell, j \leq D$) and note that $z \in D_2^1$. Similarly as in the proof of [Lemma 7.1](#) we find that $|D_i^i \cap \Gamma_{i-1}(z)|$ is independent of x, y, z , and that $|\Gamma_{i-1}(v) \cap D_2^1| = c_i$ for $v \in D_i^i$. Counting the number of pairs (w, v) such that $w \in D_2^1, v \in D_i^i, \partial(w, v) = i - 1$ in two different ways and using the above comments we obtain $|D_i^i \cap \Gamma_{i-1}(z)| |D_2^1| = |D_i^i| c_i$. The result now follows by [Lemma 4.2\(i\)](#) and [Lemma 4.3](#). ■

Corollary 7.4. *With reference to [Notation 3.1](#), for $1 \leq i \leq D$ we have*

$$E_1^* A_{i-1} E_i^* A_i E_1^* = a_i c_i k_i k^{-1} b_1^{-1} E_1^* A_2 E_1^*.$$

Proof. For $z, w \in X$ we show that the (z, w) -entries of both sides are equal. If $z \notin \Gamma(x)$ or $w \notin \Gamma(x)$ then the (z, w) -entry of each side is 0. If $z, w \in \Gamma(x)$ then the (z, w) -entries of both sides are equal by [Lemmas 5.1, 5.4](#) and [7.3](#). The result follows. ■

Lemma 7.5. *With reference to [Notation 3.1](#), for $y, z \in \Gamma(x)$ and for $1 \leq i \leq D$ the number $|\Gamma_i(x) \cap \Gamma_i(y) \cap \Gamma_i(z)|$ is equal to $a_i k_i k^{-1}$ if $y = z$ and $a_i(a_i - 1)k_i k^{-1} b_1^{-1}$ if $y \neq z$.*

Proof. Assume that $y \neq z$; otherwise the result follows by Lemma 4.3(ii). Use the abbreviation $D_j^\ell = D_j^\ell(x, y)$ ($0 \leq \ell, j \leq D$) and note that $z \in D_2^1$. It follows from Theorem 4.5 that the number of paths of length i between w and D_i^i is independent of x, y, z . Note that between any two vertices of Γ which are at distance $i - 1$ (resp. i), there exist exactly $c_1 c_2 \cdots c_{i-1} (a_2 + a_3 + \cdots + a_{i-1})$ (resp. $c_1 c_2 \cdots c_i$) paths of length i . By Lemma 7.3 the number $|D_i^i \cap \Gamma_{i-1}(z)|$ is independent of x, y, z . Using these comments we find that $|D_i^i \cap \Gamma_i(z)|$ is independent of x, y, z . For $v \in D_i^i$ we have $|\Gamma_i(v) \cap \Gamma(x)| = a_i$ so $|\Gamma_i(v) \cap D_2^1| = a_i - 1$. Counting the number of pairs (w, v) such that $w \in D_2^1, v \in D_i^i, \partial(w, v) = i$ in two different ways and using the above comments we obtain $|D_i^i \cap \Gamma_i(z)| |D_2^1| = |D_i^i| (a_i - 1)$. The result now follows by Lemmas 4.2(i) and 4.3. ■

Corollary 7.6. *With reference to Notation 3.1, for $1 \leq i \leq D$ we have*

$$E_1^* A_i E_i^* A_i E_1^* = a_i k_i k^{-1} E_1^* + a_i (a_i - 1) k_i k^{-1} b_1^{-1} E_1^* A_2 E_1^*.$$

Proof. For $z, w \in X$ we show that the (z, w) -entries of both sides are equal. If $z \notin \Gamma(x)$ or $w \notin \Gamma(x)$ then the (z, w) -entry of each side is 0. If $z, w \in \Gamma(x)$ then the (z, w) -entries of both sides are equal by Lemmas 5.1, 5.4 and 7.5. The result follows. ■

8. The irreducible T-modules with endpoint 1

With reference to Notation 3.1, in this section we describe the irreducible T -modules with endpoint 1.

Lemma 8.1. *With reference to Notation 3.1 let W denote an irreducible T -module with endpoint 1. Then $JW = 0$.*

Proof. Since W is not the primary module we have $E_0 W = 0$. Recall that $J = |X| E_0$ so $JW = 0$. ■

Lemma 8.2. *With reference to Notation 3.1 let W denote an irreducible T -module with endpoint 1. Then the following (i)–(iii) hold for $w \in E_1^* W$.*

- (i) $E_1^* A w = 0$.
- (ii) $E_i^* A_{i-1} w + E_i^* A_i w + E_i^* A_{i+1} w = 0$ for $1 \leq i \leq D - 1$.
- (iii) $E_D^* A_{D-1} w + E_D^* A_D w = 0$.

Proof. For each equation in Corollary 5.3 apply both sides to w and simplify using $E_1^* w = w$ and Lemma 8.1. ■

Lemma 8.3. *With reference to Notation 3.1 let W denote an irreducible T -module with endpoint 1. Then the following (i)–(iv) hold for $w \in E_1^* W$ and $1 \leq i \leq D$.*

- (i) $\|E_i^* A_{i-1} w\|^2 = c_i (k - c_i) k_i k^{-1} b_1^{-1} \|w\|^2$.
- (ii) $\langle E_i^* A_{i-1} w, E_i^* A_i w \rangle = -a_i c_i k_i k^{-1} b_1^{-1} \|w\|^2$.
- (iii) $\langle E_i^* A_i w, E_i^* A_{i-1} w \rangle = -a_i c_i k_i k^{-1} b_1^{-1} \|w\|^2$.
- (iv) $\|E_i^* A_i w\|^2 = a_i (k - a_i) k_i k^{-1} b_1^{-1} \|w\|^2$.

Proof. (i) Evaluating $\|E_i^* A_{i-1} w\|^2$ using $E_1^* w = w$, line (1), and Corollary 7.2 we find

$$\|E_i^* A_{i-1} w\|^2 = \frac{c_i k_i}{k} \|w\|^2 + \frac{c_i (c_i - 1) k_i}{k b_1} \langle w, E_1^* A_2 w \rangle. \quad (6)$$

By Lemma 8.2(i),(ii) we have $E_1^* A_2 w = -w$. The result follows from this and (6).

(ii), (iii), (iv) Similar to the proof of (i) above. ■

Lemma 8.4. With reference to Notation 3.1 let W denote an irreducible T -module with endpoint 1. Then for $w \in E_1^* W$ and $2 \leq i \leq D - 1$ the determinant of

$$\begin{pmatrix} \|E_i^* A_{i-1} w\|^2 & \langle E_i^* A_{i-1} w, E_i^* A_i w \rangle \\ \langle E_i^* A_i w, E_i^* A_{i-1} w \rangle & \|E_i^* A_i w\|^2 \end{pmatrix}$$

is equal to

$$a_i b_i c_i k_i^2 k^{-1} b_1^{-2} \|w\|^4.$$

Proof. Evaluate the matrix entries using Lemma 8.3 and take the determinant. ■

Theorem 8.5. With reference to Notation 3.1 let W denote an irreducible T -module with endpoint 1 and fix a nonzero $w \in E_1^* W$. Then the following is a basis for W :

$$E_i^* A_{i-1} w, \quad E_{i+1}^* A_{i+1} w \quad (1 \leq i \leq D - 1). \quad (7)$$

Proof. We first show that W is spanned by the vectors (7). Let W' denote the subspace of V spanned by the vectors (7) and note that $W' \subseteq W$. We claim that W' is a T -module. By construction W' is M^* -invariant. It follows from Lemmas 6.2–6.4 and 8.2 and $E_1^* w = w$ that W' is invariant under each of L, F, R . Recall that $L + F + R = A$ and A generates M so W' is M -invariant. The claim follows. Note that $W' \neq 0$ since $w \in W'$ so $W' = W$ by the irreducibility of W .

We now show that the vectors (7) are linearly independent. Because of (5) it suffices to show: $w \neq 0$, the vectors $E_i^* A_{i-1} w, E_i^* A_i w$ are linearly independent for $2 \leq i \leq D - 1$, and $E_D^* A_D w \neq 0$. First note that $w \neq 0$ by assumption and $E_D^* A_D w \neq 0$ by Lemma 8.3(iv). For $2 \leq i \leq D - 1$ the vectors $E_i^* A_{i-1} w, E_i^* A_i w$ are linearly independent since their matrix of inner products has nonzero determinant by Lemma 8.4. ■

We emphasize an idea from the proof of Theorem 8.5.

Corollary 8.6. With reference to Notation 3.1 let W denote an irreducible T -module with endpoint 1. Then the following (i)–(iii) hold for a nonzero $w \in E_1^* W$.

(i) $E_1^* W$ has a basis

$$w.$$

(ii) For $2 \leq i \leq D - 1$ the subspace $E_i^* W$ has a basis

$$E_i^* A_{i-1} w, \quad E_i^* A_i w.$$

(iii) $E_D^* W$ has a basis

$$E_D^* A_D w.$$

Remark 8.7. With reference to [Notation 3.1](#) let W denote an irreducible T -module with endpoint 1 and fix a nonzero $w \in E_1^*W$. It follows from [Lemma 8.3\(ii\)](#) that for $2 \leq i \leq D-1$ the basis for E_i^*W given in [Corollary 8.6\(ii\)](#) is not orthogonal. However, using the Gram–Schmidt orthogonalization one finds that $E_i^*A_{i-1}w, E_i^*A_{i-1}w + (k - c_i)a_i^{-1}E_i^*A_iw$ is an orthogonal basis for E_i^*W .

Corollary 8.8. With reference to [Notation 3.1](#) let W denote an irreducible T -module with endpoint 1. Then the following (i)–(iii) hold.

- (i) E_1^*W has dimension 1.
- (ii) E_i^*W has dimension 2 for $2 \leq i \leq D-1$.
- (iii) E_D^*W has dimension 1.

Proof. Immediate from [Corollary 8.6](#). ■

Corollary 8.9. With reference to [Notation 3.1](#) let W denote an irreducible T -module with endpoint 1. Then the following (i), (ii) hold.

- (i) The dimension of W is $2D-2$.
- (ii) The diameter of W is $D-1$.

Proof. Immediate from [Corollary 8.8](#). ■

Corollary 8.10. With reference to [Notation 3.1](#) let W denote an irreducible T -module with endpoint 1 and fix a nonzero $w \in E_1^*W$. Then $W = M^*Mw$.

Proof. By construction $M^*Mw \subseteq W$ and equality holds in view of [Theorem 8.5](#). ■

9. Irreducible T -modules with endpoint 1: the A -action

With reference to [Notation 3.1](#) let W denote irreducible T -module with endpoint 1. In this section we display the action of A on the basis for W given in [Theorem 8.5](#). Since $A = L + F + R$ it suffices to give the actions of L, F, R on this basis.

Lemma 9.1. With reference to [Notation 3.1](#) let W denote an irreducible T -module with endpoint 1 and fix a nonzero $w \in E_1^*W$. Then the matrix L from [Definition 6.1](#) satisfies the following (i)–(v).

- (i) $Lw = 0$.
- (ii) $LE_2^*Aw = (k - c_2)w$.
- (iii) $LE_i^*A_{i-1}w = (b_{i-1} + c_{i-1} + \rho_i - c_i)E_{i-1}^*A_{i-2}w + (b_{i-1} + c_{i-1} + \rho_i - c_i - \tau_{i-1})E_{i-1}^*A_{i-1}w$ for $3 \leq i \leq D-1$.
- (iv) $LE_2^*A_2w = -a_2w$.
- (v) $LE_i^*A_iw = (a_{i-1} - a_i - \rho_i)E_{i-1}^*A_{i-2}w + (\tau_{i-1} + a_{i-1} - a_i - \rho_i)E_{i-1}^*A_{i-1}w$ for $3 \leq i \leq D$.

Proof. For each equation of [Lemma 6.2](#), apply each side to w and simplify using $E_1^*w = w$ and [Lemma 8.2](#). ■

Lemma 9.2. With reference to [Notation 3.1](#) let W denote an irreducible T -module with endpoint 1 and fix a nonzero $w \in E_1^*W$. Then the matrix F from [Definition 6.1](#) satisfies the following (i)–(iv).

- (i) $Fw = 0$.
- (ii) $FE_i^*A_{i-1}w = (a_{i-1} - \rho_i)E_i^*A_{i-1}w + (c_i - \sigma_i)E_i^*A_iw$ for $2 \leq i \leq D-1$.
- (iii) $FE_i^*A_iw = (a_i - a_{i-1} + \rho_i - \rho_{i+1})E_i^*A_{i-1}w + (a_i - b_i - c_i + \sigma_i + \tau_i - \rho_{i+1})E_i^*A_iw$ for $2 \leq i \leq D-1$.
- (iv) $FE_D^*A_Dw = (a_{D-1} - c_D + \sigma_D - \rho_D)E_D^*A_Dw$.

Proof. For each equation of Lemma 6.3, apply each side to w and simplify using $E_1^*w = w$ and Lemma 8.2. ■

Lemma 9.3. With reference to Notation 3.1 let W denote an irreducible T -module with endpoint 1 and fix a nonzero $w \in E_1^*W$. Then the matrix R from Definition 6.1 satisfies the following (i)–(v).

- (i) $RE_i^*A_{i-1}w = c_iE_{i+1}^*A_iw$ for $1 \leq i \leq D-2$.
- (ii) $RE_{D-1}^*A_{D-2}w = -c_{D-1}E_D^*A_Dw$.
- (iii) $RE_i^*A_iw = \rho_{i+1}E_{i+1}^*A_iw + \sigma_{i+1}E_{i+1}^*A_{i+1}w$ for $2 \leq i \leq D-2$.
- (iv) $RE_{D-1}^*A_{D-1}w = (\sigma_D - \rho_D)E_D^*A_Dw$.
- (v) $RE_D^*A_Dw = 0$.

Proof. For each equation of Lemma 6.4, apply each side to w and simplify using $E_1^*w = w$ and Lemma 8.2. ■

10. The isomorphism class of irreducible T -modules with endpoint 1

With reference to Notation 3.1, in this section we prove that up to an isomorphism there exists a unique irreducible T -module with endpoint 1.

Theorem 10.1. With reference to Notation 3.1, any two irreducible T -modules with endpoint 1 are isomorphic.

Proof. Let W and W' denote irreducible T -modules with endpoint 1 and fix nonzero $w \in E_1^*W$, $w' \in E_1^*W'$. By Theorem 8.5, W and W' have bases $\{E_i^*A_{i-1}w, E_{i+1}^*A_{i+1}w \mid 1 \leq i \leq D-1\}$ and $\{E_i^*A_{i-1}w', E_{i+1}^*A_{i+1}w' \mid 1 \leq i \leq D-1\}$, respectively. Let $\sigma: W \rightarrow W'$ denote the vector space isomorphism defined by $\sigma(E_i^*A_{i-1}w) = E_i^*A_{i-1}w'$ and $\sigma(E_{i+1}^*A_{i+1}w) = E_{i+1}^*A_{i+1}w'$ for $1 \leq i \leq D-1$. We show that σ is a T -module isomorphism. Since A generates M and $E_0^*, E_1^*, \dots, E_D^*$ is a basis for M^* , it suffices to show that σ commutes with each of $A, E_0^*, E_1^*, \dots, E_D^*$.

Using the assertion (iv) below the line (4) and the definition of σ we immediately find that σ commutes with each of $E_0^*, E_1^*, \dots, E_D^*$. It follows from Lemmas 9.1–9.3 that σ commutes with each of L, F, R . Recall that $A = L + F + R$ so σ commutes with A . The result follows. ■

With reference to Notation 3.1 recall that V is an orthogonal direct sum of irreducible T -modules. Let W denote an irreducible T -module. By the multiplicity with which W appears in V we mean the number of irreducible T -modules in this sum which are isomorphic to W . For example the primary module $M\hat{x}$ appears in V with multiplicity 1.

Corollary 10.2. With reference to Notation 3.1, let W denote an irreducible T -module with endpoint 1. Then W appears in V with multiplicity $k-1$.

Proof. Write V as an orthogonal direct sum of irreducible T -modules W_0, W_1, \dots, W_ℓ . Without loss of generality we assume that $W_0 = M\hat{x}$. Note that $E_1^*V = \sum_{i=0}^\ell E_1^*W_i$ (direct sum) so

$$\dim E_1^*V = \sum_{i=0}^\ell \dim E_1^*W_i. \quad (8)$$

We examine the terms in (8). We mentioned earlier that $\dim E_1^*V = k$. By [7, Theorem 5.4] $E_1^*W_0$ is spanned by $\sum_{y \in \Gamma(x)} \hat{y}$ so $\dim E_1^*W_0 = 1$. By Corollary 8.8 and construction, for $1 \leq i \leq \ell$ the dimension of $E_1^*W_i$ is 1 (resp. 0) if the endpoint of W_i is 1 (resp. at least 2). Evaluating (8) using these comments we find that there exist exactly $k - 1$ T -modules among W_0, W_1, \dots, W_ℓ that have endpoint 1. The result follows in view of Theorem 10.1. ■

Corollary 10.3. *With reference to Notation 3.1 fix a nonzero $w \in E_1^*V$ which is orthogonal to $\sum_{y \in \Gamma(x)} \hat{y}$. Then M^*Mw is an irreducible T -module with endpoint 1.*

Proof. Let H denote the subspace of V spanned by the irreducible T -modules with endpoint 1. By construction and Lemma 8.1 E_1^*H is the orthogonal complement of $\sum_{y \in \Gamma(x)} \hat{y}$ in E_1^*V . Hence $w \in E_1^*H$. Note that $Tw \subseteq H$ so Tw is the orthogonal direct sum of some irreducible T -modules of endpoint 1. Call these T -modules W_1, W_2, \dots, W_s . Since $w \in Tw$ we have $w = w_1 + w_2 + \dots + w_s$ for some $w_i \in E_1^*W_i$ ($1 \leq i \leq s$). Without loss of generality we may assume that $w_1 \neq 0$. Since $w_1 \in Tw$ there exists $t \in T$ such that $w_1 = tw$. But now $w_1 = tw = tw_1 + tw_2 + \dots + tw_s$, forcing $tw_i = 0$ for $2 \leq i \leq s$ and $tw_1 = w_1 \neq 0$. But since W_1, W_2, \dots, W_s are mutually isomorphic by Theorem 10.1 this is a contradiction unless $w_2 = \dots = w_s = 0$. Therefore $w = w_1$ and $Tw = W_1$. It follows that Tw is an irreducible T -module with endpoint 1. The result follows since $Tw = M^*Mw$ by Corollary 8.10. ■

11. The case of classical parameters

The distance-regular graph Γ is said to have *classical parameters* (D, b, α, β) whenever the intersection numbers of Γ satisfy

$$c_i = \begin{bmatrix} i \\ 1 \end{bmatrix} \left(1 + \alpha \begin{bmatrix} i-1 \\ 1 \end{bmatrix} \right) \quad (1 \leq i \leq D), \quad (9)$$

$$b_i = \left(\begin{bmatrix} D \\ 1 \end{bmatrix} - \begin{bmatrix} i \\ 1 \end{bmatrix} \right) \left(\beta - \alpha \begin{bmatrix} i \\ 1 \end{bmatrix} \right) \quad (0 \leq i \leq D-1), \quad (10)$$

where

$$\begin{bmatrix} j \\ 1 \end{bmatrix} := 1 + b + b^2 + \dots + b^{j-1}.$$

In this case b is an integer and $b \notin \{0, -1\}$ [2, Proposition 6.2.1]. Assume that Γ has classical parameters (D, b, α, β) . By (9), (10) and $a_i = k - b_i - c_i$ we find

$$a_i = \begin{bmatrix} i \\ 1 \end{bmatrix} \left(\beta - 1 + \alpha \left(\begin{bmatrix} D \\ 1 \end{bmatrix} - \begin{bmatrix} i \\ 1 \end{bmatrix} - \begin{bmatrix} i-1 \\ 1 \end{bmatrix} \right) \right) \quad (1 \leq i \leq D). \quad (11)$$

In particular

$$a_1 = \beta - 1 + \alpha b \begin{bmatrix} D-1 \\ 1 \end{bmatrix}. \quad (12)$$

By [2, Corollary 8.4.2] Γ has a Q -polynomial structure such that

$$b(\theta_1^* - \theta_i^*) = \theta_0^* - \theta_{i-1}^* \quad (1 \leq i \leq D). \quad (13)$$

Pick an integer i ($2 \leq i \leq D$). By a *parallelogram of length i* in Γ we mean a 4-tuple $uvzw$ of vertices of Γ such that $\partial(u, v) = 1$, $\partial(z, w) = 1$, $\partial(u, z) = i$, $\partial(u, w) = i - 1$, $\partial(v, w) = i - 1$, $\partial(v, z) = i - 1$.

Theorem 11.1. *With reference to Notation 3.1 the following (i)–(iii) are equivalent.*

- (i) Γ has classical parameters (D, b, α, β) .
- (ii) $\rho_i = 0$ for $3 \leq i \leq D$, where ρ_i is from Theorem 4.5.
- (iii) Γ contains no parallelograms of length i for $3 \leq i \leq D$.

Suppose (i)–(iii) hold. Then $\beta = 1 - \alpha b \begin{bmatrix} D-1 \\ 1 \end{bmatrix}$. Moreover $b < -1$.

Proof. (i) \Rightarrow (ii) For $3 \leq i \leq D$ we evaluate the expression for ρ_i in Theorem 4.5 using (13) and routinely find $\rho_i = 0$.

(ii) \Rightarrow (iii) Assume that there exists a parallelogram of some length i ($3 \leq i \leq D$). Since x was chosen arbitrarily we may assume that this parallelogram is of the form $xyzw$ for some $y, z, w \in X$. Use the abbreviation $D_j^\ell = D_j^\ell(x, y)$ for $0 \leq j, \ell \leq D$ and note that $w \in D_{i-1}^{i-1}$, $z \in D_{i-1}^i$. But then $\rho_i \neq 0$ in view of Theorem 4.5, a contradiction.

(iii) \Rightarrow (i) This follows from [24, Theorem 1.1]. ■

Suppose (i)–(iii) hold. Then $\beta = 1 - \alpha b \begin{bmatrix} D-1 \\ 1 \end{bmatrix}$ by (12) and since $a_1 = 0$. Now $b < -1$ by [24, Lemma 3.3]. ■

Theorem 11.2. *With reference to Notation 3.1 assume that Γ has classical parameters (D, b, α, β) . Then the following (i), (ii) hold.*

- (i) The intersection numbers a_i, b_i, c_i of Γ satisfy

$$\begin{aligned} a_i &= -\frac{\alpha(b+1)(b^i-1)(b^{i-1}-1)}{(b-1)^2}, \\ b_i &= \frac{(b^D-b^i)(b-1-\alpha(b^D+b^i-b-1))}{(b-1)^2}, \\ c_i &= \frac{(b^i-1)(\alpha b^{i-1}+b-\alpha-1)}{(b-1)^2} \quad (0 \leq i \leq D). \end{aligned}$$

- (ii) The integers ρ_i, σ_i, τ_i from Theorem 4.5 satisfy

$$\rho_i = 0, \quad \sigma_i = c_{i-1} - b^{i-2}, \quad \tau_i = b_i, \quad (2 \leq i \leq D).$$

Proof. (i) Evaluate a_i, b_i, c_i using (11), (10), (9) and $\beta = 1 - \alpha b \begin{bmatrix} D-1 \\ 1 \end{bmatrix}$.

(ii) Note that $a_1 = 0$ implies $\rho_2 = 0$ and that for $3 \leq i \leq D$ we have $\rho_i = 0$ by Theorem 11.1. Evaluating the expressions for σ_i, τ_i in Theorem 4.5 using (13) we routinely find $\tau_i = b_i$ and $\sigma_i/c_i = (b^{i-2} - 1)/(b^i - 1)$. The result now follows using (i) above. ■

We finish the paper with two examples.

Example 11.3. The *Hermitian forms graph* has classical parameters (D, b, α, β) , where $b = -r$, r a prime power, $\alpha = b - 1$, $\beta = -b^D - 1$ [2, p. 194, Table 6.1]. Using (12) we find $a_1 = 0$ if and only if $r = 2$. Assume that $r = 2$. Then using Theorem 11.2 we find

$$a_i = \frac{2^{2i-1} - (-2)^{i-1} - 1}{3}, \quad b_i = \frac{4^D - 4^i}{3}, \quad c_i = \frac{(-2)^{i-1}(1 - (-2)^i)}{3}$$

for $0 \leq i \leq D$ and

$$\rho_i = 0, \quad \sigma_i = \frac{(-2)^{i-1}(1 - (-2)^{i-2})}{3}, \quad \tau_i = \frac{4^D - 4^i}{3} \quad (2 \leq i \leq D).$$

Example 11.4. The *Witt graph* M_{23} has classical parameters $(3, -2, -2, 5)$ [2, p. 194, Table 6.1]. Using Theorem 11.2 we find

$$a_1 = 0, \quad a_2 = 2, \quad a_3 = 6, \quad b_0 = 15, \quad b_1 = 14, \quad b_2 = 12, \quad c_1 = 1, \quad c_2 = 1, \quad c_3 = 9$$

and

$$\rho_2 = 0, \quad \rho_3 = 0, \quad \sigma_2 = 0, \quad \sigma_3 = 3, \quad \tau_2 = 12, \quad \tau_3 = 0.$$

Remark 11.5. Beside Examples 11.3 and 11.4 we are not aware of any Q -polynomial distance-regular graph with diameter $D \geq 3$ such that $a_1 = 0$ and $a_i \neq 0$ for $2 \leq i \leq D$.

Acknowledgements

The author would like to thank Paul Terwilliger for his careful reading of the earlier versions of this manuscript. The author gratefully acknowledges support by the US Department of State and the Fulbright Scholar Program, and thanks the University of Wisconsin-Madison for hospitality during his visit in Spring 2007. This work was supported in part by “Javna agencija za raziskovalno dejavnost Republike Slovenije”, program no. Z1-9614.

References

- [1] E. Bannai, T. Ito, Algebraic Combinatorics I: Association schemes, in: Benjamin-Cummings Lecture Notes, vol. 58, Menlo Park, 1984.
- [2] A.E. Brouwer, A.M. Cohen, A. Neumaier, Distance-Regular Graphs, Springer-Verlag, Berlin, Heidelberg, 1989.
- [3] J. Caughman, The Terwilliger algebras of bipartite P - and Q -polynomial association schemes, Discrete Math. 196 (1999) 65–95.
- [4] J. Caughman, M. MacLean, P. Terwilliger, The Terwilliger algebra of an almost-bipartite P - and Q -polynomial scheme, Discrete Math. 292 (2005) 17–44.
- [5] B. Collins, The girth of a thin distance-regular graph, Graphs Combin. 13 (1997) 21–30.
- [6] B. Collins, The Terwilliger algebra of an almost-bipartite distance-regular graph and its antipodal 2-cover, Discrete Math. 216 (2000) 35–69.
- [7] B. Curtin, Bipartite distance-regular graphs, Part I, Graphs Combin. 15 (1999) 143–158.
- [8] B. Curtin, Bipartite distance-regular graphs, Part II, Graphs Combin. 15 (1999) 377–391.
- [9] B. Curtin, The local structure of a bipartite distance-regular graph, European J. Combin. 20 (1999) 739–758.
- [10] G. Dickie, Twice Q -polynomial distance-regular graphs are thin, European J. Combin. 16 (1995) 555–560.
- [11] G. Dickie, P. Terwilliger, A note on thin P -polynomial and dual-thin Q -polynomial symmetric association schemes, J. Algebraic Combin. 7 (1998) 5–15.
- [12] E. Egge, A generalization of the Terwilliger algebra, J. Algebra 233 (2000) 213–252.

- [13] E. Egge, The generalized Terwilliger algebra and its finite-dimensional modules when $d = 2$, *J. Algebra* 250 (2002) 178–216.
- [14] J. Go, The Terwilliger algebra of the hypercube, *European J. Combin.* 23 (2002) 399–429.
- [15] J. Go, Tight distance-regular graphs and the subconstituent algebra, *European J. Combin.* 23 (2002) 793–816.
- [16] S.A. Hobart, T. Ito, The structure of nonthin irreducible T -modules of endpoint 1: Ladder bases and classical parameters, *J. Algebraic Combin.* 7 (1998) 53–75.
- [17] T. Ito, K. Tanabe, P. Terwilliger, Some algebra related to P - and Q -polynomial association schemes, in: *Codes and Association Schemes* (Piscataway NJ, 1999), Amer. Math. Soc., Providence RI, 2001, pp. 167–192.
- [18] T. Ito, P. Terwilliger, The shape of a tridiagonal pair, *J. Pure Appl. Algebra* 188 (2004) 145–160.
- [19] T. Ito, P. Terwilliger, Tridiagonal pairs and the quantum affine algebra $U_q(\widehat{sl}_2)$, *Ramanujan J.* 13 (2007) 41–64.
- [20] T. Ito, P. Terwilliger, Two non-nilpotent linear transformations that satisfy the cubic q -Serre relations, *J. Algebra Appl.* 6 (2007) 477–503.
- [21] Š Miklavič, Q -polynomial distance-regular graphs with $a_1 = 0$, *European J. Combin.* 25 (2004) 911–920.
- [22] K. Nomura, A refinement of the split decomposition of a tridiagonal pair, *Linear Algebra Appl.* 403 (2005) 1–23.
- [23] K. Nomura, P. Terwilliger, The split decomposition of a tridiagonal pair, *Linear Algebra Appl.* 424 (2007) 339–345.
- [24] Y. Pan, M. Lu, C. Weng, Triangle-free Distance-regular Graphs, *J. Algebraic Combin.* 27 (2008) 23–34.
- [25] K. Tanabe, The irreducible modules of the Terwilliger algebras of Doob schemes, *J. Algebraic Combin.* 6 (1997) 173–195.
- [26] P. Terwilliger, The subconstituent algebra of an association scheme, *J. Algebraic Combin.* Part I 1 (1992) 363–388. Part II: 2 (1993), 73–103; Part III: 2 (1993), 177–210.
- [27] M. Tomiyama, N. Yamazaki, The subconstituent algebra of a strongly regular graph, *Kyushu J. Mat.* 48 (1998) 323–334.